arXiv:1401.4296v3 [math.NT] 25 Nov 2014

VALUES OF TWISTED ARTIN L-FUNCTIONS

KENNETH WARD

Dedicated to the memory of Paul J. Sally, Jr.

ABSTRACT. This note gives a simple proof that certain values of Artin's L-function, for a representation ρ with character χ_ρ, are stable under twisting by an even Dirichlet character χ, up to the $\dim(\rho)$th power of the Gauss sum $\tau(\chi)$ and an element generated over $\mathbb{Q}$ by the values of χ and χ_ρ. This extends a result due to J. Coates and S. Lichtenbaum.

1. INTRODUCTION

We let $K|F$ denote a finite Galois extension of algebraic number fields, and χ will denote the character associated with a representation of $\mathrm{Gal}(K|F)$. Via C.L. Siegel and H. Klingen [4, 8], if χ denotes a Dirichlet character and K is a totally real number field, then the Dirichlet L-function $L(\chi,s)$ lies in $\mathbb{Q}(\chi)$ when s takes negative integer values. For a representation ρ, the associated character will simply be written as χ_ρ. For general representations ρ when K is a finite Galois extension of $\mathbb{Q}$, J. Coates and S. Lichtenbaum [1] decomposed the factors at infinity in the functional equation [5, VII.12.6]

$$\Lambda(K|F,\chi_\rho,s) = W(\chi_\rho)\Lambda(K|F,\overline{\chi}_\rho,1-s) \qquad (|W(\chi_\rho)|=1) \tag{1}$$

to show that, at a negative integer $s=m$ that is a critical point for the Artin L-function $L(K|\mathbb{Q},\chi_\rho,s)$, either

(I) m is odd, and the fixed field K_ρ of the kernel of ρ is totally real; or
(II) K_ρ is totally imaginary, conjugation is central in $\mathrm{Gal}(K_\rho|\mathbb{Q})$, and $\chi_\rho(\sigma) = -\dim(\rho)$.

They then employed Brauer's theorem on induced characters and functorial properties of Artin's L-function to prove that $L(K|\mathbb{Q},\chi_\rho,m)\in\mathbb{Q}(\chi_\rho)$ when m is a negative integer and critical point of $L(K|\mathbb{Q},\chi_\rho,s)$ (for all other negative integers m, this L-function trivially takes the value zero).

We consider a finite Galois extension $K|\mathbb{Q}$, and a representation χ_ρ of $\mathrm{Gal}(K|\mathbb{Q})$. We let χ denote an even Dirichlet character, which is viewed as either a one-dimensional character of $\mathrm{Gal}(K|\mathbb{Q})$ acting trivially on the conjugation automorphism, or any one-dimensional character of $\mathrm{Gal}(K|\mathbb{Q})$ if K is totally real. We also let $\mathbb{Q}(\chi_\rho,\chi)$ denote the field generated over $\mathbb{Q}$ by the values of χ_ρ and χ. The

1991 *Mathematics Subject Classification.* 11F67, 11F80, 11L05, 11M06.

Key words and phrases. Artin L-function, twisted, Dirichlet character, Gauss sum, special values.

character $\chi \otimes \chi_\rho = \chi \cdot \chi_\rho$ is associated with the representation $\chi \otimes \rho = \chi \cdot \rho$. With $\mathfrak{f} = \mathfrak{f}(\chi)$ the Artin conductor and K_χ the fixed field of the kernel of χ, we recognize the character χ as a character of the ray class group $J_{\mathbb{Q}}^{\mathfrak{f}}/P_{\mathbb{Q}}^{\mathfrak{f}}$ via the Artin symbol $(K_\chi|\mathbb{Q},\cdot) : J_{\mathbb{Q}}^{\mathfrak{f}}/P_{\mathbb{Q}}^{\mathfrak{f}} \to \mathrm{Gal}(K_\chi|\mathbb{Q})$. The Gauss sum of χ is defined in the usual way: For an algebraic number field F with ring of integers $\mathfrak{o}_F$, a Dirichlet character χ of F with finite part χ_f as a Größencharakter and conductor $\mathfrak{f}$, $\mathfrak{d}_{F|\mathbb{Q}}$ the different of $F|\mathbb{Q}$, $y \in \mathfrak{f}^{-1}\mathfrak{d}_{F|\mathbb{Q}}^{-1}$, and $Tr_{F|\mathbb{Q}}(\cdot)$ the trace from F to $\mathbb{Q}$, the Gauss sum of χ at y is given by

$$\tau(\chi,y) = \sum_{\substack{x \bmod \mathfrak{f} \\ (x,\mathfrak{f})=1}} \chi_f(x) e^{2\pi i Tr_{F|\mathbb{Q}}(xy)},$$

where the sum is taken over a set of representatives of $(\mathfrak{o}_F/\mathfrak{f})^*$. We also define χ to be zero for any ideals not relatively prime to $\mathfrak{f}$ and set $\tau(\chi) = \tau(\chi,1)$. Finally, for a field K, a subfield $F \subset K$, and two elements $a,b \in K$, we will say that $a \sim_F b$ if $a = kb$ for some $k \in F$. We may now state our main result.

Theorem 1. *If $s = m$ is a negative integer and critical point of $L(K|\mathbb{Q},\chi_\rho,s)$, then*

$$L(K|\mathbb{Q},\chi\otimes\chi_\rho,1-m) \sim_{\mathbb{Q}(\chi_\rho,\chi)} \tau(\chi)^{\dim(\rho)} L(K|\mathbb{Q},\chi_\rho,1-m).$$

Viewing χ_ρ as a character of $\mathrm{Gal}(K_\rho|\mathbb{Q})$, we may write, by Brauer's theorem on induced characters [2, 5.2],

$$\chi_\rho = \sum_{i=1}^{l} n_i \mathrm{Ind}_{H_i}^{G}(\chi_i), \tag{2}$$

where $G = \mathrm{Gal}(K_\rho|\mathbb{Q})$, and for each $i = 1,\ldots,l$, χ_i is a character of degree one of a subgroup $H_i \leq G$ and $n_i \in \mathbb{Z}$. By Galois theory, for each $i = 1,\ldots,l$, let F_i be the subfield of K_ρ so that $H_i = \mathrm{Gal}(K_\rho|F_i)$. We also let $G_i = \mathrm{Gal}(K|F_i)$ for each $i = 1,\ldots,l$. The notation in the decomposition (2) will be used henceforth. We first relate the Gauss sums of restrictions of χ to L-function values. We will show that if $s = m$ is a negative integer and critical point of $L(K|\mathbb{Q},\chi_\rho,s)$, then

$$\prod_{i=1}^{l} (\tau(\chi|_{G_i}))^{n_i} \sim_{\mathbb{Q}(\chi_\rho,\chi)} \frac{L(K|\mathbb{Q},\overline{\chi}_\rho,1-m)}{L(K|\mathbb{Q},\overline{\chi\otimes\chi}_\rho,1-m)}. \tag{3}$$

We will also show that the product of restricted Gauss sums is also related to the value of the unrestricted Gauss sum:

$$\tau(\chi)^{\dim(\rho)} \sim_{\mathbb{Q}(\chi)} \prod_{i=1}^{l} (\tau(\chi|_{G_i}))^{n_i}. \tag{4}$$

It is no surprise that ramification groups have a role in this, as we must measure the action of characters on primes. The importance of requiring that χ be an even Dirichlet character is clear: this ensures that the critical points of $L(K|\mathbb{Q},\chi\otimes\chi_\rho,s)$ and $L(K|\mathbb{Q},\chi_\rho,s)$ are the same, by conditions (I) and (II).

2. The Main Result

Our arguments take advantage of the interaction of Brauer's theorem with the Artin L-function as in the original proof of its meromorphicity. We shall also employ ideal numbers, Galois theory, and local behavior of characters. Let us turn immediately to the proof of Theorem 1.

Proof of Theorem 1. Let us fix $i \in \{1,...,l\}$. We may choose ideal numbers d_i, f_i of F_i so that the different $\mathfrak{d}_{F_i|\mathbb{Q}}$ of $F_i|\mathbb{Q}$ satisfies $\mathfrak{d}_{F_i|\mathbb{Q}} = (d_i)$ and the conductor $\mathfrak{f}_i = \mathfrak{f}(\chi_i)$ of χ_i satisfies $\mathfrak{f}_i = (f_i)$ [5, VII.7]. We let $p_i = (p_{i,\tau})_{\tau \in \mathrm{Hom}(F_i,\mathbb{C})}$ be the exponent of the infinite part of the decomposition of χ_i as a Größencharakter,

$$\chi_i((x)) = \chi_{i,f}(x)\chi_{i,\infty}(x) = \chi_{i,f}(x) N_{F_i|\mathbb{Q}}\left(\left(\frac{x}{|x|}\right)^{p_i}\right),$$

where $N_{F_i|\mathbb{Q}}(\cdot)$ denotes the norm from F_i to $\mathbb{Q}$. As χ_i is a character of the group of ideals of F_i via the Artin map, it may be viewed as a Dirichlet character, and thus has $p_{i,\tau} = 0$ unless $\tau \in \mathrm{Hom}(F_i,\mathbb{R})$. The root number $W(\chi_i)$ from the functional equation (1) satisfies

$$\begin{aligned} W(\chi_i) &= \left[\omega_4^{Tr_{F_i|\mathbb{Q}}(\overline{p}_i)} N_{F_i|\mathbb{Q}}\left(\left(\frac{f_i d_i}{|f_i d_i|}\right)^{\overline{p}_i}\right)\right]^{-1} \frac{\tau(\chi_i)}{\sqrt{\mathscr{N}_{F_i|\mathbb{Q}}(\mathfrak{f}_i)}} \\ &= \pm\left[\omega_4^{Tr_{F_i|\mathbb{Q}}(\overline{p}_i)}\right]^{-1} \frac{\tau(\chi_i)}{\sqrt{\mathscr{N}_{F_i|\mathbb{Q}}(\mathfrak{f}_i)}}, \end{aligned}$$

where $Tr_{F_i|\mathbb{Q}}(\cdot)$ denotes the trace from F_i to $\mathbb{Q}$, $\mathscr{N}_{F_i|\mathbb{Q}}(\cdot)$ the ideal norm from F_i to $\mathbb{Q}$, and ω_4 the complex number i, to avoid confusion with indices. We let d_{F_i} denote the discriminant of F_i, and we let

$$N_{i,+} = \frac{[F_i:\mathbb{Q}]}{2} + \frac{1}{2}\sum_{\substack{\mathfrak{p}|\infty \\ \mathfrak{p}\text{ real}}} \chi_i(\phi_{\mathfrak{P}}), \qquad N_{i,-} = \frac{[F_i:\mathbb{Q}]}{2} - \frac{1}{2}\sum_{\substack{\mathfrak{p}|\infty \\ \mathfrak{p}\text{ real}}} \chi_i(\phi_{\mathfrak{P}}),$$

where for each prime $\mathfrak{p}$ appearing in these sums, $\mathfrak{P}$ may be any choice of prime of K so that $\mathfrak{P}|\mathfrak{p}$ and $\phi_{\mathfrak{P}}$ is a generator of the decomposition group $G_{\mathfrak{P}}(K|F_i)$. This yields the following expression for the functional equation:

(5)

$$\begin{aligned} L(K|F_i,\chi_i,1-s) = \pm\left[\omega_4^{Tr_{F_i|\mathbb{Q}}(\overline{p}_i)}\right]^{-1} \frac{\tau(\chi_i)}{\sqrt{\mathscr{N}_{F_i|\mathbb{Q}}(\mathfrak{f}_i)}} [|d_{F_i}|\mathscr{N}_{F_i|\mathbb{Q}}(\mathfrak{f}_i)]^{s-\frac{1}{2}} \left(\cos\left(\frac{\pi s}{2}\right)\right)^{N_{i,+}} \\ \times \left(\sin\left(\frac{\pi s}{2}\right)\right)^{N_{i,-}} (2(2\pi)^{-s}\Gamma(s))^{[F_i:\mathbb{Q}]} L(K|F_i,\overline{\chi}_i,s). \end{aligned}$$

All of this holds equally well with $\chi|_{G_i} \otimes \chi_i$ in place of χ_i. For an automorphism $\sigma \in \mathrm{Gal}(\overline{\mathbb{Q}}|\mathbb{Q}(\chi_\rho,\chi))$, induction by characters (2) and functoriality of the Artin

L-function give

$$\frac{L(K|\mathbb{Q},\chi\otimes\chi_\rho,m)}{L(K|\mathbb{Q},\chi_\rho,m)}=\alpha\left[\prod_{i=1}^{l}\left(\frac{\tau(\chi|_{G_i}\otimes\sigma(\chi_i))}{\tau(\sigma(\chi_i))}\right)^{n_i}\right]\frac{L(K|\mathbb{Q},\overline{\chi\otimes\chi_\rho},1-m)}{L(K|\mathbb{Q},\overline{\chi_\rho},1-m)}, \tag{6}$$

where $\alpha\in\mathbb{Q}$. The automorphism σ holds the following fixed:

(1.a) the (rational-valued) infinite part of either Größencharakter $\chi|_{G_i}\otimes\chi_i$ or χ_i;
(1.b) the conductors of $\chi|_{G_i}\otimes\chi_i$ or χ_i; and
(1.c) the action of $\chi|_{G_i}\otimes\chi_i$ or χ_i on the generators of decomposition groups for infinite primes.

By the functional equation (5) without the use of σ, it follows from (1.a)-(1.c) that

$$\frac{L(K|\mathbb{Q},\chi\otimes\chi_\rho,m)}{L(K|\mathbb{Q},\chi_\rho,m)}=\alpha\left[\prod_{i=1}^{l}\left(\frac{\tau(\chi|_{G_i}\otimes\chi_i)}{\tau(\chi_i)}\right)^{n_i}\right]\frac{L(K|\mathbb{Q},\overline{\chi\otimes\chi_\rho},1-m)}{L(K|\mathbb{Q},\overline{\chi_\rho},1-m)}, \tag{7}$$

so that (6) and (7) give

$$\prod_{i=1}^{l}\left(\frac{\tau(\chi|_{G_i}\otimes\sigma(\chi_i))}{\tau(\sigma(\chi_i))}\right)^{n_i}=\prod_{i=1}^{l}\left(\frac{\tau(\chi|_{G_i}\otimes\chi_i)}{\tau(\chi_i)}\right)^{n_i}. \tag{8}$$

Furthermore, as in Theorem 1.2 of [1], we have

$$\frac{L(K|\mathbb{Q},\chi\otimes\chi_\rho,m)}{L(K|\mathbb{Q},\chi_\rho,m)}\in\mathbb{Q}(\chi_\rho,\chi), \tag{9}$$

by none other than the fact that the value of each L-function in the numerator and denominator of (9) lies in $\mathbb{Q}(\chi_\rho,\chi)$. We let ζ be a root of unity, chosen so that $\mathbb{Q}(\zeta)$ contains all of the values of the characters χ and χ_i, as well as any roots of unity appearing in the Gauss sums $\tau(\chi_i)$, $\tau(\chi|_{G_i}\otimes\chi_i)$, and $\tau(\chi|_{G_i})$, for all $i=1,\ldots,l$. Let $\sigma\in\mathrm{Gal}(\mathbb{Q}(\zeta)|\mathbb{Q}(\chi_\rho,\chi))$, so that $\sigma(\zeta)=\zeta^d$ for some integer d. We apply σ to the expression

$$\prod_{i=1}^{l}\left(\frac{\tau(\chi|_{G_i}\otimes\chi_i)}{\tau(\chi|_{G_i})\,\tau(\chi_i)}\right)^{n_i}. \tag{10}$$

We note that, for any Dirichlet character χ of an algebraic number field, $\tau(\chi,a)=\overline{\chi}_f(a)\tau(\chi)$ if $(a,\mathfrak{f})=1$, but also that $\tau(\chi,a)=0=\overline{\chi}_f(a)\tau(\chi)$ if $(a,\mathfrak{f})\neq 1$, as we have defined χ to be zero for ideals not relatively prime to $\mathfrak{f}$. Thus, for each $i=1,\ldots,l$,

(2.a)
$$\begin{aligned}\sigma(\tau(\chi|_{G_i}\otimes\chi_i))&=\tau(\chi|_{G_i}\otimes\sigma(\chi_i),d)\\&=\left(\overline{(\chi|_{G_i}\otimes\sigma(\chi_i))}_f(d)\right)\tau(\chi|_{G_i}\otimes\sigma(\chi_i));\end{aligned}$$

(2.b) $\sigma(\tau(\chi|_{G_i}))=\tau(\sigma(\chi|_{G_i}),d)=\tau(\chi|_{G_i},d)=\left(\overline{(\chi|_{G_i})}_f(d)\right)\tau(\chi|_{G_i})$; and

(2.c) $\sigma(\tau(\chi_i))=\tau(\sigma(\chi_i),d)=\left(\overline{\sigma(\chi_i)}_f(d)\right)\tau(\sigma(\chi_i))$.

This requires that d be relatively prime to the conductors of each of these characters. By (8) and (2.a)-(2.c), we find that σ fixes (10). By (6), (9), we obtain (3).

As in the previous argument, we again select an appropriate choice of root of unity ζ, which in this instance contains the values of the characters $\chi|_{G_i}$ and χ, as well as all roots of unity appearing in $\tau(\chi|_{G_i})$ and $\tau(\chi)$, and we consider some $\sigma \in \mathrm{Gal}(\mathbb{Q}(\zeta)|\mathbb{Q}(\chi))$, where $\sigma(\zeta) = \zeta^d$ for some integer d. As before, d must be relatively prime to the conductors of χ and $\chi|_{G_i}$, for all $i = 1,...,l$. Similarly to previous arguments, we obtain

$$\sigma\left(\frac{\prod_{i=1}^{l}(\tau(\chi|_{G_i}))^{n_i}}{\tau(\chi)^{\dim(\rho)}}\right) = \left(\frac{\prod_{i=1}^{l}\left(\overline{(\chi|_{G_i})}_f(d)\right)^{n_i}}{\overline{\chi}_f(d)^{\dim(\rho)}}\right)\left(\frac{\prod_{i=1}^{l}(\tau(\chi|_{G_i}))^{n_i}}{\tau(\chi)^{\dim(\rho)}}\right). \tag{11}$$

Note that σ makes no appearance in this final expression, as σ fixes the values of χ and $\chi|_{G_i}$, for all $i = 1,...,l$. Let K_χ and $K_{\chi|_{G_i}}$ denote the fixed fields of the kernels of χ and $\chi|_{G_i}$, for each $i = 1,...,l$, respectively. We have for each $i = 1,...,l$ that $(\chi|_{G_i})_f(d) = \chi|_{G_i}((d))$, where (d) is viewed as an ideal of $K_{\chi|_{G_i}}$, and likewise that $\chi_f(d) = \chi((d))$, where (d) is viewed as an ideal of K_χ. We write $d = \prod_k p_k^{a_k}$ as the prime factorization of d in $\mathbb{Z}$. As d is relatively prime to the conductor of χ, each prime p_k is unramified in K_χ. Thus, for a prime $\mathfrak{P}|\mathfrak{p}_k$ of K with decomposition group $G_\mathfrak{P}$ and inertia group $I_\mathfrak{P}$ of $\mathfrak{P}|\mathfrak{p}_k$, the character χ may be viewed as a character of $G_\mathfrak{P}/I_\mathfrak{P}$.

Let us fix some $i \in \{1,...,l\}$. For each k, we write $p_k = \prod_j \mathfrak{P}_{k,j}^{e_{\mathfrak{P}_{k,j}}}$ in K and $p_k = \prod_m \mathfrak{P}'^{e_{\mathfrak{P}'_{k,m}}}_{k,m}$ in F_i. For each j, we write $\phi_{\mathfrak{P}_{k,j}}$ for a representative element of $G_{\mathfrak{P}_{k,j}}$ which maps to the Frobenius in the residue field of $\mathfrak{P}_{k,j}|\mathfrak{p}_k$, and we write $\phi'_{\mathfrak{P}_{k,j}}$ for the analogous representative in $\mathrm{Gal}(K|F_i)$ [5, VII.10]. Let us also fix a choice of k and j. Let $\mathfrak{P}'_{k,m}$ denote the prime of F_i with $\mathfrak{P}_{k,j}|\mathfrak{P}'_{k,m}|\mathfrak{p}_k$, and let $f_{\mathfrak{P}'_{k,m}} = [\mathfrak{o}_{F_i}/\mathfrak{P}' : \mathbb{Z}/p_k\mathbb{Z}]$. By definition of $\phi_{\mathfrak{P}_{k,j}}$ and $\phi'_{\mathfrak{P}_{k,j}}$, the image

$$\overline{\phi_{\mathfrak{P}_{k,j}}^{f_{\mathfrak{P}'_{k,m}}}}(\alpha \bmod \mathfrak{P}_{k,j}) = \phi_{\mathfrak{P}_{k,j}}^{f_{\mathfrak{P}'_{k,m}}}(\alpha) \bmod \mathfrak{P}_{k,j}$$

in the residue field of $\mathfrak{P}_{k,j}|\mathfrak{p}_k$ is equal to $\phi'^{f_{\mathfrak{P}'_{k,m}}}_{\mathfrak{P}_{k,j}}(\alpha) \bmod \mathfrak{P}_{k,j}$. Thus $\phi_{\mathfrak{P}_{k,j}}^{f_{\mathfrak{P}'_{k,m}}}$ and $\phi'_{\mathfrak{P}_{k,j}}$ belong to the same coset of $I_{\mathfrak{P}_{k,j}}$ in $G_{\mathfrak{P}_{k,j}}$. As χ is trivial on $I_{\mathfrak{P}_{k,j}}$, we have $\chi(\phi_{\mathfrak{P}_{k,j}}^{f_{\mathfrak{P}'_{k,m}}}) = \chi(\phi'_{\mathfrak{P}_{k,j}})$. By the identity $\sum_m e_{\mathfrak{P}'_{k,m}} f_{\mathfrak{P}'_{k,m}} = [F_i : \mathbb{Q}]$, we obtain $\chi((d))^{[F_i:\mathbb{Q}]} = \chi|_{G_i}((d))$. Returning to Brauer's theorem on induced characters (2), we also have $\dim(\rho) = \sum_{i=1}^{l} n_i[F_i : \mathbb{Q}]$. It follows that

$$\prod_{i=1}^{l}\left((\chi|_{G_i})_f(d)\right)^{n_i} = \chi_f(d)^{\dim(\rho)}. \tag{12}$$

By (11) and (12), σ fixes $\prod_{i=1}^{l}(\tau(\chi|_{G_i}))^{n_i}/\tau(\chi)^{\dim(\rho)}$, which must therefore lie in $\mathbb{Q}(\chi)$. This proves (4). Finally, as $\tau(\chi)\tau(\overline{\chi}) \in \mathbb{Q}(\chi)$, we obtain from (3) and (4)

that

$$L(K|\mathbb{Q},\chi\otimes\chi_\rho,1-m)\sim_{\mathbb{Q}(\chi_\rho,\chi)}\tau(\chi)^{\dim(\rho)}L(K|\mathbb{Q},\chi_\rho,1-m),$$

proving Theorem 1. □

3. Acknowledgements

The author would like to thank New York University for its support while this paper was written, as well as Anantharam Raghuram for posing this question on Gauss sums and L-functions.

References

[1] J. Coates and S. Lichtenbaum. On l-adic zeta functions. *Annals of Mathematics*. **98** (1973), 498-550.
[2] M. Collins. *Representations and Characters of Finite Groups*. Cambridge, 1990.
[3] E. Freitag and R. Busam. *Complex Analysis*. Springer-Verlag, 2005.
[4] H. Klingen. Über die Werte der Dedekindschen Zetafunktion. *Mathematische Annalen*. **145** (1961/2), 265-272.
[5] J. Neukirch. *Algebraic Number Theory*. Springer-Verlag, 1999.
[6] J.-P. Serre. *Linear Representations of Finite Groups*. Springer-Verlag, 1977.
[7] T. Shintani. On evaluation of zeta functions of totally real algebraic number fields at non-positive integers. *Journal of the Faculty of Science, University of Tokyo. Section 1A, Mathematics*. **2 (23)** (1976), 393-417.
[8] C.L. Siegel. Über die analytische Theorie der quadratischen Formen. III. *Annals of Mathematics*. **1 (2) 38** (1937), 212-291.

NYU-ECNU Institute of Mathematical Sciences, New York University Shanghai, Shanghai, China, 200062
E-mail address: kaw354@nyu.edu